



DSSSB TGT

PART (A+B)



MATHS

COMPLEX ANALYSIS

PART-10



20-09-2024 04:00 PM



Complex Analysis

Algebraic Number: - A complex number is said to be Algebraic number if it satisfies a non-zero polynomial over $\mathbb{Q}$.

Transcendental Number: - Non -algebraic numbers are said to be transcendental number.

Q. Which of the following numbers is an algebraic number?

A. π

B. e

C. $\sqrt{2}$

D. $\ln(2)$

Q. Which of the following statements is true?

A, All transcendental numbers are algebraic.

B. All algebraic numbers are transcendental.

C. Some algebraic numbers are transcendental.

D. No algebraic number is transcendental.

Complex Analysis

(1) $|z - a| = r$

(2) $|z - a| < r$

(3) $|z - a| \leq r$

(4) $|z - a| > r$

(5) $|z - a| \geq r$

Complex Analysis

Analytic Function or Regular Function or
Holomorphic Function

Analytic Function:- A complex valued function $f(z)$ is said to be analytic at point $z = z_0$ if $f'(z) = \lim_{\delta z \rightarrow 0} \frac{f(z+\delta z) - f(z)}{\delta z}$ exists & is unique at z_0

Analytic Function: A single valued function $f(z)$ is said to be analytic in a region R , if it is differentiable at each point of R .

(C-R-Equation | Cauchy Riemann Equation)

For a function $f(z)$ to be analytic in a region R

Necessary: If $f(z)$ is analytic at a point, then u and v must satisfy the Cauchy-Riemann equations at that point.

Sufficient: If u and v satisfy the Cauchy-Riemann equations and the partial derivatives of u and v are continuous, then $f(z)$ is analytic in that region

Q. Let $f(z) = \frac{1}{z}, z \neq 0$, then

- 1. f does not satisfy Cauchy - Riemann equation for all $z \neq 0$**
- 2. f is continuous only but nowhere differentiable**
- 3. f satisfies Cauchy - Riemann equation, but not analytic for all $z \neq 0$**
- 4. f is analytic for all $z \neq 0$,**

Q. The Complex Function $f(z) = z^2$ is?

- (a) Analytic**
- (b) Not analytic**
- (c) May be analytic**
- (d) None of these**

Q. $f(z) = (|z|)^2$ is analytic or not?

- (a) Analytic**
- (b) Not analytic**
- (c) May be analytic**
- (d) None**

Q. $f(z) = 2xy + i(x^2 - y^2)$

- (a) Analytic**
- (b) Not analytic**
- (c) May be analytic**
- (d) None**

Q. $f(z) = z^3$ $(x+iy)^3$

☒ (a) Analytic

☐ (b) Not analytic

☐ (c) May be analytic

☐ (d) None

$$x^3 + i^3 y^3 + 3x^2 iy + 3xi^2 y^2$$

$$x^3 - iy^3 + 3x^2 yi - 3xy^2$$

$$\underbrace{(x^3 - 3xy^2)}_u + i \underbrace{(3x^2 y - y^3)}_v$$

Q. $f(z) = e^{-x}(\cos y + i \sin y)$

- (a) Analytic**
- (b) Not analytic**
- (c) May be analytic**
- (d) None**

Q. $f(z) = \sin z$

- (a) Analytic**
- (b) Not analytic**
- (c) May be analytic**
- (d) None**

Harmonic Function

Q. $u = x^2 - y^2$

- (a) Harmonic function
- (b) Not Harmonic function
- (c) May be Harmonic function
- (d) None

Characteristics of Harmonic Functions in Complex Analysis

- 1.If $f(z) = u(x, y) + iv(x, y)$ is analytic in a region R , then both u and v are harmonic functions in R .
- 2.If $u(x, y)$ is harmonic in a connected region R , then u is the real part of an analytic function $f(z) = \underbrace{u(x, y)} + i\underbrace{v(x, y)}$.
- 3.If u and v are the real and imaginary parts of an analytic function, then u and v are considered as harmonic conjugates.

⇒ 4. The sum of two harmonic functions results in another harmonic function.

5. An arbitrary pair of harmonic functions " u " and " v " may not necessarily be conjugates, unless $u + iv$ is an analytic function.

Q. $v = 3x^2y - y^3$

- ✓ (a) Harmonic function
- (b) Not Harmonic function
- (c) May be Harmonic function
- (d) None

$$u_{xx} + u_{yy} = 0$$

$$v = 3x^2y - y^3$$

$$v_x = 6xy$$

$$v_{xx} = 6y$$

$$V = 3x^2y - y^3$$

$$V_y = 3x^2 - 3y^2$$

$$V_{yy} = -6y$$

$$v_{xx} + v_{yy} = 0$$

For analytic fun $f(z) = u + iv$

Q. $u(x, y) = x^2 - y^2$, find $v(x, y)$?

C-R Eqn

$$u_x = v_y$$

$$u_y = -v_x$$

~~(a) $3x + 2y + k$~~

(b) $2xy + k \Rightarrow v_y = 2x$

(c) $3xy^2 + k \Rightarrow v_y = 6xy$

~~(d) None~~

$$u_x = 2x = v_y$$

$$u = x^2 - y^2$$
$$u_y = -2y$$

$$v = 2xy + k$$
$$v_x = 2y$$

For an analytic fun
 $\Downarrow f(z) = u + iv$

C-R Eqⁿ

$$\begin{aligned} u_x &= v_y \\ u_y &= -v_x \end{aligned}$$

$$\begin{aligned} u &= x^3 - 3xy^2 + c \\ u_y &= -6xy \end{aligned}$$

Q. $v(x, y) = 3x^2y - y^3$, Find $u(x, y)$

~~(a)~~ $x^2 - 3xy^3 + c$

~~(b)~~ $x^3 - 2xy + c$

(c) $x^3 - 3xy^2 + c$

(d) None

$$u_x = 3x^2 - 3y^2$$

$$v = 3x^2y - y^3$$

$$v_y = 3x^2 - 3y^2$$

$$v = 3x^2y - y^3$$

$$v_x = 6xy$$

Q. For analytic function $f(z) = u + iv$, **if** $u(x, y) = 3x^2 - 3y^2$, **find** $v(x, y)$

~~(a)~~ $6xy + 3x^2 + c$

(b) $6y - 6x + c$

(c) $6x - 6y + c$

(a) $6xy + c$

$$v_x = 6y + 6x^2$$

$$v = 6xy + c$$

$$v_x = 6y$$

C-R Eqⁿ

$$u_x = v_y$$

$$u_y = -v_x$$

$$u = 3x^2 - 3y^2$$

$$u_x = 6x \checkmark$$

$$u_y = -6y$$

C-R Eqⁿ

$$\begin{aligned} u_x &= v_y \\ u_y &= -v_x \end{aligned}$$

Q. An analytic Function $f(z) = u(x, y) + iv(x, y)$ **if** $u = xy$, **find** v

~~(a)~~ $\frac{x^2 - y}{2} + k$

~~(b)~~ $\frac{x - y^2}{2} + k$

~~(c)~~ $\frac{(x+y)^2}{2} + k$

(d) $\frac{y^2 - x^2}{2} + k$

$$\begin{aligned} u &= xy \\ u_x &= y = v_y = y \\ u_y &= x = -v_x \end{aligned}$$

$$v = \frac{y^2 - x^2}{2} + k$$

$$v_x = \frac{1}{2}(-2x) = -x$$

Milne -Thomson Method

Construct Analytic function By Milne -Thomson Method

(1) $f(z) = \int (u_x)_{(z,0)} dz - i \int (u_y)_{(z,0)} dz + c$; **if** $u(x, y)$ **is given**

(2) $f(z) = \int (v_y)_{(z,0)} dz + i \int (v_x)_{(z,0)} dz + c$; **if** $v(x, y)$ **is given**



Q. If $u(x, y) = x^3 - 3xy^2 + 3x^2 - 3y^2$, **Find Analytic**
 $f(z)$?

$$3x^2 - 3y^2 + 6x$$

(a) $z^3 - z^2 + c$

(b) $z^3 + 3z^2 + c$

(c) $3z^3 + z^2 + c$

(d) None of these.

$$f(z) = \int (u_x)_{(z,0)} dz - i \int (u_y)_{(z,0)} dz + c$$

$$\int (3z^2 - 3(0)^2 + 6z) dz - i \int 0 dz + c$$

$$\rightarrow \frac{3z^3}{3} + \frac{6z^2}{2} + c \Rightarrow z^3 + 3z^2 + c$$

Q. $u(x, y) = x^3 - 3xy^2$. Find Analytic $f(z)$?

(a) $z^2 + c$ $3x^2 - 3y^2$

(b) $z^3 + z^2 + c$

(c) $z^3 + c$

(d) None of these.

$-6xy$

$$f(z) = \int (U_x)_{(z,0)} dz - i \int (U_y)_{(z,0)} dz + C$$

$$\int 3z^2 dz - i \int 0 dz + C$$

$$\frac{3z^3}{3} + C$$

$$f(z) = z^3 + C$$

Q. $V(x, y) = 3x^2y - y^3$. Find Analytic $f(z)$?

(a) $z^2 + c$ $3x^2 - 3y^2$

(b) $z^3 + z^2 + c$ $6xy$

(c) $z^3 + c$

(d) None of these.

$$f(z) = \int (V_y)_{(z,0)} dz + i \int (V_x)_{(z,0)} dz + C$$

$$\Rightarrow \int 3z^2 dz + i \int 0 dz + C$$

$$\Rightarrow \frac{3z^3}{3} + C = z^3 + C$$